

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

FENDER 240 INSECTICIDE

ACTIVE CONSTITUENT: 240g/L METHOXYFENOZIDE
ALSO CONTAINS: 60g/L ETHYLENE GLYCOL
1.5g/L PARAFORMALDEHYDE

GROUP **18** INSECTICIDE

For the control of Lightbrown Apple Moth on Apples and Pears (Pome Fruit), Blueberries, Citrus, Grapevines and Kiwi Fruit; and for the control of various other lepidopteran pests of Almonds, Apples and Pears (Pome Fruit), Avocado, Coffee, Custard Apple, Eggplant, Longan, Lychee, Peppers (Capsicum and Chilli), Okra, Macadamia and Tomatoes as specified in the Directions For Use table.

APVMA Approval No.: 95188/144839

Pack Size: 250mL-1000L



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | titanag.com.au

SC

Formulation Type
Suspension Concentrate

IN A TRANSPORT EMERGENCY
• DIAL 000 • POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING NOT
A DANGEROUS GOOD ACCORDING
TO THE AUSTRALIAN DANGEROUS
GOODS (ADG) CODE FOR TRANSPORT
BY ROAD AND RAIL

DIRECTIONS FOR USE

Restrictions:

DO NOT apply if rain or overhead irrigation is expected within 6 hours. Retain the first flush of irrigation tailwater/storm water in the tailwater dam after application.

DO NOT use more than 3 sprays per season in any single crop.

DO NOT use in covered or protected situations such as glasshouses, greenhouses, shade houses or plastic tunnels.

DO NOT apply by aircraft.

DO NOT apply more than 3 times to any Almond crop.

DO NOT apply more than once at the hull split stage for Almond crops.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings 1 to 2 hours before sunset and persist until 1 to 2 hours after sunrise.

TREE AND VINE CROPS			
In the following table, all rates given are for dilute spraying. For concentrate spraying, refer to the MIXING/APPLICATION section. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. For concentrate spraying, do NOT use at rates greater than 5 times the dilute spraying rate.			
CROP	PEST	RATE/100L	CRITICAL COMMENTS
Almonds	Carob Moth (1st generation)	80mL + Wetter	Practice good orchard hygiene with the removal of mummified fruit in winter. Early post-flower application: At the beginning of the season (Aug-Sept), carefully monitor pheromone traps for peak adult pest incidence of 1st generation. Once regular field scouting indicates egg laying and newly hatched larvae, apply a maximum of 3 sprays of TITAN Fender 240 Insecticide at a minimum of 10 day intervals between sprays. Apply to complete coverage. Refer to Wetting Agents in the General Instructions section for more information.
	Carob Moth (2nd generation)		Hull-split application: TITAN Fender 240 Insecticide should be applied at the start of hull split (2-5% of the Almond hull sutures open). Carefully monitor pheromone traps for peak activity of 2nd generation moths. Continue field scouting for egg-laying activity and apply TITAN Fender 240 Insecticide to coincide with the initiation of egg hatch. Apply to ensure complete coverage of all foliage and fruit (hull/shell) surfaces. Concentrate spraying is not appropriate for this pest. DO NOT apply less than 2000L/ha water.
Apples and Pears (and other Pome Fruit)	Lightbrown Apple Moth	25mL	For control of the spring generation of Lightbrown Apple Moth, commence spraying at petal fall and apply in a series of 3 applications at 14 day intervals. Alternatively, in areas where a later summer generation occurs, spray when moth activity is indicated by pheromone traps or lure pots. Apply a minimum of 2 sprays with a 21 day interval. Ensure thorough coverage of developing fruit clusters. When insect pressure is high in blocks with large, vase-shaped trees and with short-stemmed, bunchy varieties, TITAN Fender 240 Insecticide must be applied by dilute spraying to ensure run-off into fruit clusters.
	Loopers		Control of these pests is best achieved using a schedule of 3 sprays of TITAN Fender 240 Insecticide in spring (see above).
Avocado	Avocado Leafroller		Crop monitoring is recommended from pre-bloom. Apply at first sign of pest incidence and target eggs and newly hatched larvae. Additional sprays may be required if re-infestation occurs.
Blueberry	Lightbrown Apple Moth		Thorough coverage of the plant is essential. Target eggs and newly hatched larvae. Apply when locally determined thresholds are exceeded.
Citrus		25mL + a wetting agent at 10mL/100L	Use with high volume spraying equipment only. Thorough coverage is essential. Spray to run-off. Apply when eggs and very small larvae are first seen in flower clusters or developing fruitlets. A second spray may be required 2-3 weeks later if larvae hatch over an extended period.
Coffee	Avocado Leafroller	25mL	Crop monitoring for pest incidence is recommended from pre-bloom. Apply at first sign of pest incidence, targeting eggs and very small larvae before they are protected in retreats. Additional sprays may be required if re-infestation occurs.
Custard Apple	Yellow Peach Moth		Apply when locally determined pest thresholds are exceeded as indicated by field checks. Additional sprays may be required if re-infestation occurs. Spray to thoroughly cover fruit.
Grapevines	Lightbrown Apple Moth	25mL	Thorough leaf tip and bunch coverage is essential. Spring generation: Apply from pre-flowering onwards targeting eggs and small (5-6mm long) larvae (up to 3rd instar). Apply a second spray 14-21 days later. Summer generation: Apply at first sign of activity targeting eggs and larvae up to 5-6mm long (up to 3rd instar). The summer generation can only be detected by monitoring.
Kiwi Fruit			Monitor pest activity from bud burst onwards. Target sprays at eggs and first instar larvae.

CROP	PEST	RATE/100L	CRITICAL COMMENTS
Longan	Macadamia Nut Borer	40mL	Spray thoroughly when pest numbers reach economic threshold levels according to field checks. Target sprays against eggs and early instar larvae.
Lychee			
Macadamia	Macadamia Flower Caterpillar	25mL	Monitor for eggs and very small larvae on flowers and apply at a threshold of 50-80% of racemes infested.
	Macadamia Nut Borer	40mL	Spray to thoroughly cover nuts when pest numbers reach economic threshold levels according to field checks. Target sprays against eggs and early instar larvae.
NON-TREE AND VINE CROPS			
Note: TITAN Fender 240 Insecticide is a moult-accelerating insecticide that requires ingestion for control. Feeding ceases almost immediately after ingestion. Larvae in protected feeding sites (eg flowers) will not be controlled. Mortality of larvae will not be evident until 4-6 days after application.			
CROP	PEST	RATE	CRITICAL COMMENTS
Tomatoes, Peppers (Capsicum and Chilli), Eggplant, Okra	Native Budworm, Tomato Grub, Cluster Caterpillar	125 or 170mL/100L or 1.25 or 1.7L/ha	Use the higher rate under heavy egg pressure. Apply TITAN Fender 240 Insecticide to brown eggs or at egg hatch when pest numbers reach treatment threshold levels as determined by field checks. Maintain field checks and reapply after 7 days if necessary. Ensure thorough coverage of plants.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

HARVEST

Tomato, Peppers (Capsicum and Chilli), Eggplant and Okra: NOT REQUIRED WHEN USED AS DIRECTED.

Citrus Fruits: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

Longan: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

Almond: DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

Blueberry: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

Avocado, Coffee, Custard Apple, Kiwi Fruit, Lychee and Pome Fruit: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

Grapes (for domestic consumption): DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

Grapes (for export or wine production): Consult your winery, industry spray diary or peak industry body for the recommended withholding period (export harvest interval) required to meet export residue requirements.

Macadamia: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.

GRAZING

All Crops, Orchards, Plantations and Vineyards: DO NOT GRAZE TREATED CROPS.

DO NOT ALLOW LIVESTOCK TO GRAZE WITHIN TREATED ORCHARDS, PLANTATIONS OR VINEYARDS.

Grapes, Blueberries and Coffee: DO NOT APPLY WHEN THERE ARE LIVESTOCK, PASTURE OR ANY LAND THAT IS PRODUCING FEED FOR LIVESTOCK

WITHIN 5m DOWNWIND OF THE APPLICATION AREA.

Tomato, Peppers, Eggplant and Okra: DO NOT APPLY WHEN THERE ARE LIVESTOCK, PASTURE OR ANY LAND THAT IS PRODUCING FEED FOR LIVESTOCK WITHIN 15m DOWNWIND OF THE APPLICATION AREA.

NOTE – CROPS INTENDED FOR EXPORT: TITAN Fender 240 Insecticide may leave detectable chemical residues in harvested produce. Overseas markets may not have appropriate residue tolerances in place or may have established tolerances which are lower than Australian maximum residue limits. Some crops for export to these destinations may require a longer harvest withholding period. If you plan to use this product on crops destined for export, please check with your exporter before using TITAN Fender 240 Insecticide.

GENERAL INSTRUCTIONS

TITAN Fender 240 Insecticide is very suitable for use in orchards and vineyards using integrated pest management (IPM) programs.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of TITAN Fender 240 Insecticide to water in the spray vat while stirring or with agitators in motion.

WETTING AGENTS

Not all surfactants or crop oils are of equal quality and if intending to use other wetting agents it is recommended that you consult TITAN AG Pty Ltd prior to use. TITAN AG only supports the use of Agral Spray adjuvant and Nufarm Chemwet 1000 with this formulation.

DILUTE SPRAYING

Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions For Use table for each 100L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

CONCENTRATE SPRAYING

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. This is needed to calculate the

concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500L/ha
2. Your chosen concentrate spray volume: For example 500L/ha
3. The concentration factor in this example is 3X (ie $1500L \div 500L = 3$)
4. If the dilute label rate is 25mL/100L, then the concentrate rate becomes 3 x 25, that is 75mL/100L of concentrate spray.

The chosen spray volume and the sprayer set up and operation may need to be changed as the crop grows.

DO NOT use a concentrate rate higher than that specified in the Critical Comments. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow Industry Best Practices.

COMPATIBILITY

TITAN Fender 240 Insecticide is incompatible with mineral spray oils. DO NOT mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of TITAN AG Pty Ltd, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, TITAN AG cannot be responsible for the behaviour of such mixtures.

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management TITAN Fender 240 Insecticide is a Group 18 Insecticide. Some naturally occurring insect biotypes resistant to TITAN Fender 240 Insecticide and other Group 18 Insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if TITAN Fender 240 Insecticide and other Group 18 Insecticides are used repeatedly. The effectiveness of

GROUP 18 INSECTICIDE



TITAN Fender 240 Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of TITAN Fender 240 Insecticide to control resistant insects. TITAN Fender 240 Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, TITAN AG representative or local agricultural department agronomist.

PRECAUTIONS

DO NOT apply using spray equipment carried on the back of the user.

Re-entry Period: DO NOT allow entry into treated areas until the spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied on plants at any time. Keep livestock out of operational areas during treatment. Avoid spray drift onto adjoining properties or stock areas.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic life with long lasting effects. DO NOT spray across open bodies of water. DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses. DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertilisers or seed. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For drumMUSTER containers: This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. DO NOT dispose of undiluted chemical on-site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced but may be taken separately.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to designated collection point for refill or storage.

110L Containers: If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. DO NOT attempt to breach the valve system of the filling point or contaminate the container with water or other products. Ensure that the couple, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of TITAN AG Pty Ltd.

100L, 200L and 1000L Containers: If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. The container must be vented before discharging contents. To empty, connect a camlock fitted hose to the bottom valve. Remove top cap when discharging for venting purposes. When the container is empty, close all caps and valves and return the container to the point of purchase.

SPILL AND LEAK MANAGEMENT

DO NOT touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways and drains.

Small spills/leaks: Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to TITAN AG Pty Ltd.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact, inhaled or swallowed. Attacks eyes. The fumes first cause smarting, then watering of the eyes. This should be taken as a warning sign. Will irritate nose and throat and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes, skin and clothing. DO NOT inhale vapour or spray mist. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves, goggles and half facepiece respirator with canister specified for formaldehyde. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing and respirator (and if rubber wash with detergent and warm water).

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. Remove from contaminated area. Apply artificial respiration if not breathing. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Fender 240 Insecticide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: None.
